

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Claims


Sub C1

1. A method of treating a ^{patient} ~~patent~~ for depression comprising the step of administering to said patient an effective amount of a compound having a NMDA IC₅₀ of about 50 nM to about 1 μM as measured in the NMDA assay and a serotonin reuptake IC₅₀ of less than or equal to about 100 nM as measured in the serotonin reuptake inhibition assay.

2. The method of claim 1, wherein said compound has an NMDA receptor IC₅₀ of 50 nM to 1 μM and a SSRI IC₅₀ less than 100 nM.

526
B2

3. A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



15

wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, and -O-acyl;

20

Ar¹ and Ar² are each independently selected from the group consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, cyclohexyl, cycloheptyl, and cyclopentyl;

25

each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²'s together are imino;

132 each R^1 is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl; and

each m is independently an integer from 0 to 5;

provided that if both R_3 's are $-CH_3$, then both X_m 's are not 3-F, 4-F, 3- CF_3 , 4-Cl, and if both R_3 's are $-CH_3$ and one X_m is 4-F then the other X_m is not 4-Cl; further provided that if one R_3 is -H and the other R_3 is $-CH_3$, then both X_m 's are not 4-Cl, and if one R_3 is -H and the other R_3 is $-CH_3$, then at least one m is 1; or a pharmaceutically acceptable salt thereof.

10

4. The method of claim 3 wherein for said compound each X is independently either -F, -Cl, - OCF_3 , or - CF_3 ;

each R^1 is -H;


each R^2 is -H;

15

one R^3 is -H, and the other R^3 is either -H or $-CH_3$; and

each m is 1.

5. The method of claim 3 wherein said compound has the chemical structure:



20

wherein X^1 is either -Br, -Cl, -F, -I, - CF_3 , alkyl, -OH, - OCF_3 , -O-alkyl, or -O-acyl;

X^2 is either -Br, -Cl, -F, -I, - CF_3 , alkyl, -OH, - OCF_3 , -O-alkyl, or -O-acyl; and

25

R^3 is either -H or $-CH_3$;

or a pharmaceutically acceptable salt thereof.

6. The method of claim 5, wherein X^1 is -F, -Cl, - OCF_3 , or - CF_3 ; and X^2 is either 2- OCH_3 , 2- CH_3 , 3-F, 3- CF_3 , or 4- CF_3 .


30

TOTALT-5040660

Sub C1

546
B3 ✓

7. A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



5 wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, and -O-acyl;

Ar¹ and Ar² are each independently selected from the group
 10 consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, cyclohexyl, cycloheptyl, and cyclopentyl;

each R¹ is independently selected from the group consisting
 15 of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²s together are imino;

each R³ is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl; and

20 m is 0 to 5;

or a pharmaceutically acceptable salt thereof.

8. The method of claim 7, wherein for said compound each X is independently either -F, -Cl, -OCF₃, or -CF₃;

Ar¹ and Ar² are each independently phenyl or naphthyl;

each R¹ is -H;

each R² is -H;


one R³ is -H, and the other R³ is either -H or -CH₃;

each m is 0 or 1.

00000405-112101

Sal
C1/

9. The method of claim 7, wherein said compound has the chemical structure:



wherein X¹ is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;


X² is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl; and

R³ is either -H or -CH₃;

or a pharmaceutically acceptable salt thereof.

10. The method of claim 9 wherein X¹ is either -F, -Cl, -OCF₃ or -CF₃; and X² is either 2-OCH₃, 2-CH₃, 3-F, 3-CF₃, or 4-CF₃.

11. A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, and -O-acyl;

34 each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²s together are imino;

5 each R³ is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl;

Z is either -CH₂CH₂-, -CH₂CH(CH₃)-, -CH=CH-, -O-CH₂-, -S-CH₂-, -CH₂-, -O-, or -S-; and

10 each n is independently 1 to 4; or a pharmaceutically acceptable salt thereof.


12. The compound of claim 11, wherein each X is independently either -F, -Cl, -OCF₃, or -CF₃;

each R¹ is -H;

each R² is -H;

15 one R³ is -H, and the other R³ is either -H or -CH₃; and each n is 1.

13. The method of claim 11, wherein said compound has the chemical structure:



20 wherein X¹ is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;

X² is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl; and


25 R³ is either -H or -CH₃;

or a pharmaceutically acceptable salt thereof.

09900405-112104

526
135
14. The method of claim 13 wherein X^1 is -F, -Cl, -OCF₃ or -CF₃; and X^2 is either either -F, -Cl, -OCH₃, -CH₃, -OCF₃ or -CF₃.

5 15. A method of treating a patient for depression comprising the step of administering to said patient an effect amount of a compound having the chemical structure:



10 wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, and -O-acyl; ; preferably, each X is independently either -F, -Cl, -OCF₃ or -CF₃;

Ar¹ and Ar² are each independently selected from the group
15 consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, cyclohexyl, cycloheptyl, and cyclopentyl; preferably Ar¹ and Ar² are independently naphthyl or phenyl; more preferably at least one of
20 Ar¹ and Ar² is phenyl; and more preferably, both Ar¹ and Ar² are phenyl;

Y is either -CH₂-, -O-, or -S-;

each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;
25 preferably, each R¹ is -H;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²'s together are imino; preferably each R² is -H;


each R³ is independently selected from the group consisting
30 of -H, alkyl, 2-hydroxyethyl, and alkylphenyl, preferably, each R³ is independently either -H or -CH₃; more preferably one R³ is -H, and the other R³ is either -H or -CH₃; and

0990405-112101

B5

each m is independently an integer from 0 to 5; and preferably, each m is independently 0 or 1.

16. The method of claim 15, wherein said compound has the chemical structure; Structure VIII



wherein X^1 is independently selected from the group consisting of -H, -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl; preferably, X^1 is either -F, -Cl, -OCF₃ and -CF₃;


X^2 is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl; preferably, X^2 is independently either -F, -Cl, -OCH₃, -CH₃, -OCF₃ or -CF₃; more preferably, X^2 is either 2-OCH₃, 2-CH₃, 3-F, 3-CF₃, or 4-CF₃; and

R^3 is either -H or CH₃;

or a pharmaceutically acceptable salt thereof.

09990405-112101
TOTAL-5040660

17. A compound having the chemical structure;
















or a pharmaceutically acceptable salt thereof.

09900405-112101

526
B6

18. A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



5 or a pharmaceutically acceptable salt thereof.

add
B7

00990405.11103